

Bronze Age Economies and Landscape Resources in the Kargaly Steppe (Orenburg, Russia). Remote Sensing and Palynological Data for Ancient Landscape Resources Modeling

Carlos Fernández Freire¹, Antonio Uriarte González², Juan Manuel Vicent García³ and M^a Isabel Martínez Navarrete³.

Unidad de SIG, CCHS – CSIC, Madrid, Spain; carlos.ffreire@cchs.csic.es
Laboratorio de Arqueología del Paisaje y Teledetección, Instituto de Historia, CCHS – CSIC, Madrid, Spain; antonio.uriarte@cchs.csic.es
Research Group Prehistoria Social y Económica, CCHS - CSIC, Madrid, Spain; {juan.vicent, isabel.martinez}@cchs.csic.es

Abstract. We present the methodological and technical issues of a research on ancient vegetation modeling in the Russian steppes combining remote sensing and palynological data. This research is framed in an international project developed by Russian and Spanish archaeologists from the Russian Academy of Sciences and Spanish National Research Council (CSIC) and centered on the study of the Bronze Age mining complex of Kargaly (Orenburg, Russia).

One of the research guidelines focused on potential landscape resources for copper metallurgy and subsistence activities, such as the availability of wood fuel or pasture and arable land. A palaeobotanical research has been carried out in order to explore such questions, including the use of remote sensing data for providing environmental calibrative criteria that help interpreting palynological data. Combining current pollen rain data and remote sensing products (obtained by digital processing of Landsat and ASTER imagery), we have developed a model to improve our understanding of the paleopalynological record.

Keywords. Paleopalynology, Remote Sensing, Bronze Age, Prehistoric Mining, Landscape Archaeology, Eurasian Steppe.

1. Introduction

Kargaly is a vast copper mineral field located in the Orenburg region, in the forest steppe of the middle Ural river basin (Fig. 1).

First mining exploitations go back to Early Bronze Age (EBA), in the middle of the 4th millennium BC, although mining and metallurgical works reach their maximum intensity in Late Bronze Age (LBA). Since the end of this period, mining activity completely stops and does not resume till Modern Ages, with the annexation of this region by the Russian Empire in 1750 AD, and fully decays in the beginning of the 20th century.

Since 1993, the prehistoric mining-metallurgical complex of Kargaly has been the subject of research by a Russian-Spanish team managed by Dr. E. N. Chernykh (Institute of Archaeology, Russian Academy of Sciences, Moscow) and formed by researchers from the Russian Academy of Sciences, the Spanish National Research Council (CSIC) and the National Archaeological Museum in Madrid. During the course of research, several mining districts have been surveyed and mapped and three Bronze Age settlements and several Kurgan cemeteries have been excavated [1-5]. At the same time, paleoenvironmental and landscape archaeology research has been carried out, which frames the work shown in this paper.




Figure 1: Kargaly location in the Eurasian steppe close to the Ural river

The study of Kargaly landscape and its transformations since Bronze Age until present time is fundamental for the historical understanding of the Kargaly complex, its development through Bronze Age and its collapse at the end of this period. According to their relevance, two aspects are emphasized:

1. Mining and metallurgy are activities that have a great impact on environment, mainly in relation to forest resources. Changes in the distribution of these resources between Bronze Age and present time might constitute a good proxy to evaluate the scale of mining works and the existence or not of a local metallurgy.
2. Archaeological data show that agriculture was not practiced in this region till a very later period [6]. This raises some questions about the subsistence economy of the Bronze Age mining communities that settled the area.

Our research has focused on the first topic up to now [7] being subsistence practices the subject of current works.

Available data about vegetation distribution in Bronze Age are very limited and restricted to evidence provided by fossil pollen taken from archaeological sediments and natural deposits excavated during fieldwork. In order to interpret this information according to the aims of our project, we have proposed a methodology based on palynological spectra, taken by subsoil drill sampling, versus modern pollen rain patterns comparison, taken by surface soil samples [8]. Understanding the relationship between these patterns and modern vegetation, will help us interpret observable differences between fossil and recent spectra as signs of changes in the vegetation cover. This requires a precise cartography about vegetation formations in the area, which is the aim of the remote sensing program shown in this paper.

Proposed methodology is based on the *Best Modern Analogue* approach [9-11], developed by archaeological palynology in recent years. This approach consists of isolating present palynological

spectra that are qualitative and quantitatively similar to fossil pollen spectra and analyzing the environment of the first assuming a similar environment for the last. Incorporating remote sensing to the BMA methodology provides a powerful tool for representing and analyzing the environment. In this research we try to promote a synergy between palynology and remote sensing by means of statistical models.

Next, we briefly describe the general research design and the works carried out for the elaboration of a land cover cartography through satellite image processing. Finally, we discuss some of the results of the analysis of the palynological data in relation to the problem of changes in composition and distribution of forest resources between Bronze Age and present time.

2. Methods

2.1. Research design:

In order to feed the analytical landscape model that will be used for testing the paleoenvironmental hypotheses, two basic sources of information have been used: palynological spectra and land cover cartography.

Palynological spectra are elaborated through the study of palynological samples taken at different locations in the research area. They provide information about botanical taxa existing in the surroundings of each sample point. Each taxon is represented by a variable containing its relative frequency (expressed through percentage) with regard to the whole sum of pollen grains in the sample.

Two types of palynological samples have been analyzed: fossil samples and present pollen rain samples. Fossil paleopalynological samples are taken from ancient natural deposits and provide information about past vegetation. They are chrono-referenced through relative or numeric dating methods (e.g. radiocarbon). Modern pollen rain samples are taken on the ground surface level and provide information about modern episodes of pollen deposition. In order to offer a complete view of pollen rain in the whole study area, a representative set of samples must be taken. This has been achieved by means of a sampling strategy, combining both a selective and a systematic approach.

Paleopalynological samples have been taken at two natural deposits (22 samples at Novenki peat and 17 at Gorny mining trench) [12], [13]. These samples have been dated through the radiocarbon method, applied both to pollen grain concentrations and paleosols. Chronology ranges since 4th millennium BC to present time.

Pollen rain data have been obtained by means of 76 surface samples, 55 of them resulting from a systematic sampling strategy and the other 21 selectively located according to botanical criteria [7], [14] (Fig. 2). Systematic sampling was based on two adjoining transects, each one defined around one of the paleopalynological samples. Each transect was divided into square sampling units 500 m on a side. A 10% of these units was randomly selected in each transect. In each selected unit a sample was taken in the upper 10 cm of the topsoil.




Figure 2: Pollen rain and palynological samples distribution

Regarding land cover cartography, its function consists of being compared statistically with pollen rain data for exploring how and to what degree these represent the quantitative presence and spatial distribution of different botanical taxa through landscape. Due to the lack of enough detailed pre-existing cartography, it has been generated using remote sensing imagery combined with ancillary information, such as a DEM, by means of digital image classification methods. Image classification has been supervised using existing knowledge about vegetation in Kargaly region, both botanical literature[15], [16] and fieldwork observations.

The process of information analysis has the following stages (Fig. 3):

1. Quantification of the presence of pollen taxa in each sample, through the elaboration of relative frequency variables.
2. Elaboration of land cover cartography using image classification methods. Primary data are remote sensing images and ancillary data, such as a DEM.
3. Comparative analysis of pollen rain and land cover cartography by means of statistical methods. This allows to elaborate a model of how pollen rain represents vegetation presence.
4. Finally, comparison of fossil pollen data with this model, for estimating which is the most likely kind of vegetation landscape that these data represent.




Figure 3: Data integration and analysis work flow

2.2. Remote sensing:

Current landscape was to be modeled through remote sensing techniques¹, in order to obtain an analytical document displaying the vegetation distribution responsible for the pollen rain samples gathered during fieldwork campaigns.

To achieve a correct land cover classification of the study area we had to rely on very heterogeneous data. A Landsat TM 5 image accounted for terrain conditions at the moment of field data



Figure 4: Landsat TM image. False color composition

¹ ERDAS Imagine was used for digital image processing, and ESRI ArcGIS software for ancillary data such as DEM.




Figure 5: ASTER image. False color composition

gathering, but 30 m spatial resolution revealed too poor for the characterization of certain pollen species dispersal, and inadequate to delineate the narrow forest spots typical in Kargaly. Furthermore, there is no cartography of the area at an appropriate scale, so an ASTER image was acquired to enable the building of a DEM, as well as to improve spatial resolution, via data merging.

Although the Landsat scene was collected on September 1994 and the ASTER one on August 2002, data merging was considered feasible thanks to the continuities that link both landscapes. In spite of ownership and crop changes, land subdivision remain practically unchanged since soviet agrarian colonization.

The ASTER image was acquired at Level 1B, which means that it had been already geometrically corrected via orbital parameters; the TM had been geometrically corrected using accurate GCPs, consistent with pollen data samples. The ASTER image had to be co-registered to the TM image, a process that yielded a RMSE of 11 meters.

The Landsat image was merged with a panchromatic ASTER generated as an average of the three VNIR bands. We chose a method based on wavelet analysis, for it produces a higher spectral quality than other methods as PCA or HIS [17], a crucial point in order to perform a digital classification. The merged image was considered suitable according to the spectral correlation of the new image bands and the original one, as well as the ERGAS coefficient [18] (Table 1).

A supervised classification was performed on the new merged image in order to identify three main land cover categories, subdivided according to the spectral characteristics of the vegetation in autumn:

1. Forest
 - 1.1. Headwaters forest
 - 1.2. River forest
 - 1.3. Scrubland
2. Steppe

| Table 1 | |
|--------------------|-------------------|
| Band | Correlation coef. |
| 1 | 0,97435174 |
| 2 | 0,97058677 |
| 3 | 0,97316609 |
| 4 | 0,9751107 |
| 5 | 0,97536803 |
| 6 | 0,97275086 |
| | |
| ERGAS ² | 1,57232994 |

² The merging process does not alter significantly an image when the ERGAS coefficient is below 3.

- 2.1. Dry
- 2.2. Wet
- 3. Labor field
 - 3.1. Plowed fields
 - 3.2. Stubble
 - 3.3. Fallow
 - 3.4. Not harvested

A parallelepiped classification was performed with Mahalanobis distance used as overlapping rule. Afterwards forests were reclassified in river forests and headwaters forests, regarding their inclusion in first order water stream basins, according to Shreve's method³ [19]. A majority filter was applied inside parcels, so every pixel was assigned the most frequent value in its labor field, a process that required land subdivision digitizing.

Finally, the resulting image was evaluated using 62 ground control points gathered during field work. The special conditions of the field campaigns, that took place along three different years, prevented us from using all cover categories, so the image had to be recoded to the three main classes in order to perform the error matrix (Tables 2 – 4).

Table 2: Error Matrix

| Classified Data | Forest | Steppe | Labor fields |
|-----------------|--------|--------|--------------|
| Forest | 4 | 1 | 0 |
| Steppe | 1 | 35 | 0 |
| Labor fields | 0 | 0 | 21 |
| Column Total | 5 | 36 | 21 |

Table 3: Accuracy

| Class Name | Reference Totals | Classified Totals | Number Correct | Producers Accuracy | Users Accuracy |
|--|------------------|-------------------|----------------|--------------------|----------------|
| Forest | 5 | 5 | 4 | 80,00% | 80,00% |
| Steppe | 36 | 36 | 35 | 97,22% | 97,22% |
| Labor fields | 21 | 21 | 21 | 100,00% | 100,00% |
| Totals | 62 | 62 | 60 | | |
| Overall Classification Accuracy = 96.77% | | | | | |

³ Hydrological features (basins, streams) were obtained through ASTER DEM processing.

Table 4: Kappa statistics

| Class name | Kappa |
|---------------|--------|
| Forest | 0,7825 |
| Steppe | 0,9338 |
| Labor fields | 1,0000 |
| Overall Kappa | 0,9404 |




Figure 5: Digital classification results

3. Results

Current Kargaly landscape is a typical forest steppe, representing the southern strip of the Great Eurasian Steppe. It is a region of gentle hills and gullies, with a predominance of herbaceous species and scarce forest patches across the water courses, either permanent or seasonal. In our current research stage, attention has been focused on these scarce forest resources, trying to determine whether their distribution during Bronze Age was significantly different to the present one.

The study of ground truth, especially through the elaboration of systematic floristic inventories, made possible the characterization of three basic associations of woody species:

1. River forest mostly formed by willows, poplars, alders and other minority species.
2. Headwater forest, composed by birch and aspens.
3. Scrubland, made up by several species of woody *Rosaceae*.

The elaboration of detailed forest cover cartography enabled the linking of these vegetation formations with landscape topography.

With regard to the present pollen rain samples, they were statistically analyzed by means of the Principal Component Analysis technique (PCA)[7], establishing an interpretative pattern based on the quantitative variation of the first four components in relation to the location of forest species. Thus we were able to define the spatial relationship between current forest distribution and its palynological correlate.

According to the logic of the BMA approach, patterns resulting from this analysis were used to interpret diachronic pollen variations in the two paleopalynological sequences taken from natural deposits. These sequences, synchronized by means of radiocarbon dating and stratigraphical criteria, have helped to establish four stages in the formation pattern of the pollen record:

1. EBA and MBA periods (4th and 3rd millennium BC): Beginning of the mining exploitation in the region.
2. LBA period, *Srubnaya* culture (mid 3rd to mid 2nd millennium BC): Maximum intensity of the prehistoric mining exploitation.
3. Nomadic settlement period (mid 2nd millennium BC – c. 1750 AD): It is a stage without mining activity nor permanent settlement.
4. Recent period (c. 1750 AD – present time): Modern mining till c. 1900 and Soviet planned agriculture, between c. 1950 and 1991.

The comparison between fossil pollen data from these phases and observed patterns of present pollen rain yields some significant results [7]. An intense deforestation associated to phases 1 and 2 has been detected in the mining area located at the north of the study area. In the south part, where there was no mining activity, we can notice an intense impact of pastoral practices. At the same time we can observe an important variation in the composition of forest resources, consisting of practical extinction of oak (*Quercus robur*) as a consequence of anthropic impact during phase 4. This fact has relevant implications about the energetic potential of Kargaly forest resources and provides a key reference for the historical interpretation of this Bronze Age mining-metallurgical complex.

4. Conclusions

Apart from the specific results that we have discussed, research put forward in this paper shows that the proposed methodological design is feasible and relevant for Landscape Archaeology. Combining remote sensing technology and palynological research in an experimental statistical framework provides a powerful tool for evaluating landscape long-term changes, giving way to a new spatial approach to paleoenvironmental research.

Acknowledgements

The authors would like to thank E. N. Chernykh and the Institute of Archaeology (Russian Academy of Sciences, Moscow) teamwork, it is thanks to them that we could take part in the Kargaly project.

We would also like to thank all the Spaniards in charge of the palynological research: Pilar López García and Antonio López Sáez, from the Instituto de Historia (CSIC, Madrid); and to I. Zavala Morencos (Universidad Politécnica, Madrid) and A. Rodríguez Alcalde (CSIC, Madrid) in

charge of the cartography and remote sensing project, along with Juan M. Vicent during the field work.

References

- [1] E. N. Chernykh, Ed., 2002. *Gorny – the Late Bronze Age settlement. Topography, lithology, stratigraphy. Household, manufacturing and sacral structures. Relative and absolute chronology*. Moscow: Languages of Slavonic culture.
- [2] E. N. Chernykh, Ed., 2002. *Geological and geographical characteristics. History of discoveries, exploitation and investigations. Archaeological sites*. Moscow: Languages of Slavonic culture.
- [3] E. N. Chernykh, Ed., 2004. *Gorny site: Archaeological materials. Mining and metallurgy technology. Archaeobiological studies*. Moscow: Languages of Slavonic culture.
- [4] E. N. Chernykh, Ed., 2007. *Kargaly: phenomenon and paradoxes of development. Kargaly in the systems of Metallurgical Provinces Hidden (sacral) life of archaic miners and metallurgists*. Moscow: Languages of Slavonic culture.
- [5] E. N. Chernykh, Ed., 2005. *Kargaly's necropolis. Kargaly population: Palaeoanthropological investigations*. Moscow: Languages of Slavonic culture.
- [6] E. N. Chernykh, E. E. Antipina, y E. J. Lebedeva, 1998. *Produktionsformen der Urgesellschaft in den Steppen Osteuropas (Ackerbau, Viehzucht, Erzgewinnung und Verhüttung)*, in *Das Karpenbecken und die Osteuropäische Steppe. Nomadenbewegungen und Kulturaustausch in den vorchristlichen Metallzeiten (4000-500 v.Chr)*, München, 233-252.
- [7] J. M. Vicent García, M. I. Martínez Navarrete, J. A. López Sáez, y I. Zabala Morencos, 2010. *Impacto medioambiental de la minería y la metalurgia del cobre durante la Edad del Bronce en Kargaly (región de Orenburgo, Rusia)*, *Trabajos de Prehistoria*, vol. 67, nº. 2, 511-544.
- [8] J. M. Vicent García, Á. Rodríguez Alcalde, J. A. López Sáez, I. de Z. Morencos, P. López García, y M. I. Martínez Navarrete, 2000. *¿Catástrofes ecológicas en la estepa? Arqueología del paisaje en el complejo minero-metalúrgico de Kargaly (región de Orenburg, Rusia)*, *Trabajos de Prehistoria*, vol. 57, nº. 1, 29 - 74
- [9] P. Tarasov, J.W. Williams, A. Andreev, T. Nakagawa, E. Bezrukova, U. Herzschuh, Y. Igarashi, S. Müller, K. Werner, Z. Zheng, 2007. *Satellite - and pollen - based quantitative woody cover reconstructions for northern Asia: Verification and application to late-Quaternary pollen data*, *Earth and planetary science letters*, vol. 264, 284-298.
- [10] J. T. Overpeck, T. Webb III, y I. C. Prentice, 1990 *Quantitative Interpretation of Fossil Pollen Spectra: Dissimilarity Coefficients and the Method of Modern Analogs*, *Quaternary Research*, vol. 23, págs. 87-108, 1985.
- [11] J. Guiot, «Methodology of the last climatic cycle reconstruction from pollen data», *Palaeogeography, Palaeoclimatology, Palaeoecology*, vol. 80, 49-69.
- [12] J. A. López Sáez, P. López García, y M. I. Martínez Navarrete, «Palynological research at the hill of Gorny», in *Kargaly II*, E. N. Chernykh, Ed. Moscow: Languages of Slavonic culture, 2002, págs. 153-165.
- [13] P. López, J. A. López Sáez, E. N. Chernykh, y P. Tarasov, 2003 *Late Holocene vegetation history and human activity shown by pollen analysis of Novienki peat bog (Kargaly region, Orenburg Oblast, Russia)*, *Vegetation History & Archaeobotany*, vol. 12, 75-82.
- [14] J. M. Vicent García, S. Ormeño Villajos, M. I. Martínez Navarrete, y J. Delgado, 2006. *The Kargaly Project: Modelling Bronze Age landscapes in the steppe*, S. Campana y M. Forte, Eds., 279-284.
- [15] A. A. Chibilév, 1999. *Geographical Atlas of the Orenburg Oblast*. Moscow: Book Publishing House of Orenburg.
- [16] Z. N. Riabinina, 1998. *Synopsis of the flora of the Orenburg Oblast*. Ekaterinburg: Department of the Urals of the Russian Academy of Sciences.
- [17] M. González-Audicana, R. García, y A. Seco, 2002. *Fusión de imágenes multiespectrales y pancromáticas: nuevas alternativas metodológicas basadas en las transformaciones wavelet discretas empleando el algoritmo de Mallat*, *Revista de Teledetección*, nº. 18, 63-73.
- [18] L. Wald, 2000. *Quality of high resolution synthesized images: is there a simple criterion?*, in *Proceedings of the third conference Fusion of Earth data: merging point measurements, raster maps and remotely sensed images*, Sophia Antipolis, France, 99-103.
- [19] R. L. Shreve, 1966. *Statistical Law of Stream Numbers*, *The Journal of Geology*, vol. 74, nº. 1, 17-37.